

Russell Kitch, MD • Jenn Grady, MD • Jeffery Neal, MD • Julie Malka, AuD North Charleston • West Ashley

Balloon Sinuplasty

Pre-procedure Instructions

Prescriptions will be given to you at prior to your procedure.

1. Start prednisone 5 days prior to procedure or otherwise directed by your surgeon.

Day of procedure, one hour prior to arrival:

- 1. Afrin-5 sprays each nostril.
- 2. Ativan (1 mg)-take one pill; bring extra pills to office.
- 3. Vicodin (5/300)- take one pill; bring extra pills to office.
- 4. Eat a light meal prior to taking medication and procedure.

Post-procedure Instructions

- 1. Light red to clear drainage from the nose is normal for 3 to 6 days following the procedure.
- 2. Start nasal sinus rinses once a day, Nettie Pot or Neil Med sinus rinse, the first day after the procedure.
- 3. Headache, sinus or nasal discomfort is common after the procedure and you should take your prescribed pain medication for this.
 - Pain meds can give you constipation, keep hydrated and use stool softeners as needed.
- 4. No bending or lifting (over 20 pounds), strenuous activity or flying until released by your physician. Sneeze with mouth open. No nose blowing.
- 5. Follow-up will be one week after procedure.

When to Call

- Excessive, bright red blood that does not stop with pressure or Afrin.
- Persistent, excessive, clear drainage after the first week of surgery. Fever over 101°F.
- Stiffness or pain in neck or decreased alertness.
- Sudden vision changes or eye swelling.